

Finite slope cyclic surgeries along toroidal Brunnian links and generalized Properties P and R

Teruhisa KADOKAMI

April 16, 2015

Abstract

Let M_λ be the λ -component Milnor link. For $\lambda \geq 3$, we determine completely when a finite slope surgery along M_λ yields a lens space including S^3 and $S^1 \times S^2$, where *finite slope surgery* implies that a surgery coefficient of every component is not ∞ . For $\lambda = 3$ (i.e. the Borromean rings), there are three infinite sequences of finite slope surgeries yielding lens spaces. For $\lambda \geq 4$, any finite slope surgery does not yield a lens space. As a corollary, M_λ for $\lambda \geq 3$ does not yield both S^3 and $S^1 \times S^2$ by any finite slope surgery. We generalize the results for the cases of *Brunnian type links* and toroidal Brunnian type links (i.e. Brunnian type links including essential tori in the link complement). Our main tools are Alexander polynomials and Reidemeister torsions. Moreover we characterized toroidal Brunnian links and toroidal Brunnian type links in some senses.

Contents

1	Introduction	2
2	Reidemeister torsion	5
2.1	Surgery formulae	5
2.2	d -norm	7
3	Key Lemmas	8
3.1	Alexander polynomial of algebraically split links	8
3.2	Key Lemmas	10
4	Proof of Theorem 1.1	15
5	Proofs of Theorem 1.2 and Corollary 1.3	16

2010 *Mathematics Subject Classification*: 11R27, 57M25, 57M27, 57Q10, 57R65.

Keywords: Milnor link; Dehn surgery; Property P; Property R; Reidemeister torsion; Alexander polynomial.

6	Generalization to Brunnian type links and toroidal Brunnian links	19
6.1	Generalization to Brunnian type links	19
6.2	Generalization to toroidal Brunnian type links	22

1 Introduction

A λ -component link L with $\lambda \geq 1$ is a *Brunnian link* if every proper sublink of L is a trivial link. For $\lambda \geq 3$, L is algebraically split (i.e. the linking number of every 2-component sublink of L is 0), and for $\lambda \geq 1$, every component of L is an unknot. We assume that for $\lambda = 1$, L is the unknot, and for $\lambda = 2$, L is algebraically split.

Let $M_\lambda = K_1 \cup \cdots \cup K_\lambda$ be the λ -component Milnor link [Mi1, Mi2] (Figure 1). In particular, M_1 is the unknot, M_2 is the Hopf link, and M_3 is the Borromean rings. Since Dehn surgeries along the unknot and the Hopf link are well-known (for example, see [KS]), we assume $\lambda \geq 3$ for M_λ . As properties of M_λ , it is a Brunnian link, it can be constructed from the Hopf link by a sequence of Bing doubles (cf. Subsection 6.2), the components K_1 and K_2 , and $K_{\lambda-1}$ and K_λ are interchangeable, and the order of the components $(K_1, \dots, K_\lambda)$ can be reversed into $(K_\lambda, \dots, K_1)$. The Alexander polynomial of M_λ is as follows (see also Lemma 6.14 (3) and Remark 6.15) :

$$\begin{cases} \Delta_{M_3}(t_1, t_2, t_3) & \doteq (t_1 - 1)(t_2 - 1)(t_3 - 1) & (\lambda = 3), \\ \Delta_{M_\lambda}(t_1, \dots, t_\lambda) & = 0 & (\lambda \geq 4). \end{cases} \quad (1.1)$$




Figure 1: λ -component Milnor link M_λ

Let $L = K_1 \cup \cdots \cup K_\lambda$ be a λ -component link. We denote the result of $(r_1, \dots, r_\lambda)$ -surgery along L by $(L; r_1, \dots, r_\lambda)$ where $r_i \in \mathbb{Q} \cup \{\infty, \emptyset\}$ ($i = 1, \dots, \lambda$) is a surgery coefficient of K_i . A Dehn surgery producing a 3-manifold with cyclic fundamental group is called a *cyclic surgery*. In particular, a Dehn surgery producing a lens space is called a *lens surgery*. We define that an $(r_1, \dots, r_\lambda)$ -surgery along L is a *finite slope surgery* if $r_i \neq \infty, \emptyset$ for every i ($i = 1, \dots, \lambda$). When we set $r_i = p_i/q_i \in \mathbb{Q} \cup \{\infty\}$, we assume $p_i, q_i \in \mathbb{Z}$, $\gcd(p_i, q_i) = 1$ and $\pm 1/0 = \infty$. The lens space of type (p, q) is

the result of p/q -surgery along the unknot, which is denoted by $L(p, q)$. In particular, $L(1, q) \cong S^3$ and $L(0, \pm 1) \cong S^1 \times S^2$. We obtain the following theorems:

Theorem 1.1 *Let $Y = (M_3; p_1/q_1, p_2/q_2, p_3/q_3)$ be the result of a finite slope surgery along M_3 . Then Y is a lens space if and only if one of the following holds:*

- (1) $(p_1/q_1, p_2/q_2) = (\varepsilon, \varepsilon)$ and $|\varepsilon p_3 - 6q_3| = 1$, or
- (2) $(p_1/q_1, p_2/q_2) = (\varepsilon, 2\varepsilon)$ and $|\varepsilon p_3 - 4q_3| = 1$, or
- (3) $(p_1/q_1, p_2/q_2) = (\varepsilon, 3\varepsilon)$ and $|\varepsilon p_3 - 3q_3| = 1$,

and the cases that indices of (p_i, q_i) ($i = 1, 2, 3$) are permuted arbitrarily, where $\varepsilon = 1$ or -1 . Moreover, if (1), then $Y = L(p_3, 4\varepsilon q_3)$, if (2), then $Y = L(2p_3, \varepsilon(8q_3 - p_3))$, and if (3), then $Y = L(3p_3, \varepsilon(3q_3 - 2p_3))$.

Theorem 1.2 *For $\lambda \geq 4$, any finite slope surgery along M_λ does not yield a lens space.*

For a non-trivial knot K in S^3 , K has *Property P* if the fundamental group of the result of any finite slope surgery ($1/q$ -surgery with $q \neq 0$) along K is a non-trivial group, and K has *Property R* if the result of 0-surgery along K is not homeomorphic to $S^1 \times S^2$. Note that by L. Moser [Mos] which determines the results of Dehn surgery along torus knots, and by the cyclic surgery theorem due to M. Culler, M. Gordon, J. Luecke and P. Shalen [CGLS], q can be restricted to $q = \pm 1$ in the statement of Property P. We define that a non-trivial knot K in S^3 has *Property P'* if the result of any finite slope surgery along K is not S^3 , which is a stronger property than Property P. Since the Poincaré conjecture has been settled affirmatively by G. Perelman [Pe1, Pe2, Pe3], Property P and Property P' are equivalent. However, as Property P has been introduced to prove the Poincaré conjecture, we cannot use the statement of Property P' when we try to give an alternative proof of the Poincaré conjecture.

The *Property P conjecture* (*Property R conjecture*, respectively) for a knot is that a non-trivial knot in S^3 has Property P (Property R, respectively). As for the Property P conjecture for a knot, C. McA. Gordon and J. Luecke [GL] gave a great partial affirmative answer, and P. Kronheimer and T. Mrowka [KM] proved affirmatively for every non-trivial knot. As for the Property R conjecture for a knot, D. Gabai [Gb2] proved affirmatively.

We define their link versions. For a λ -component non-trivial link L in S^3 , L has *Property P* (*Property FP*, respectively) if the fundamental group of the result of any Dehn surgery (finite slope surgery, respectively) along L whose surgery coefficients are not all ∞ (i.e. $(r_1, \dots, r_\lambda) \neq (\infty, \dots, \infty)$) is non-trivial, L has *Property P'* (*Property FP'*, respectively) if the result of any Dehn surgery (finite slope surgery, respectively) along L whose surgery coefficients are not all ∞ is not S^3 , and L has *Property R* (*Property FR*, respectively) if the result of any Dehn surgery (finite slope surgery, respectively) along L is not homeomorphic to $S^1 \times S^2$. It is easy to see that not all non-trivial link

has Property P, Property FP, Property P', Property FP', Property R and Property FR even if we assume that the link is non-split. For example, the Hopf link is a such link. By the same reason as above, Property P and Property P', and Property FP and Property FP' are equivalent, but we should distinguish them for an alternative proof of the Poincaré conjecture.

A link L in S^3 has a *unique exterior* if there exists a link L' and a homeomorphism h between the complements of L and L' , then h preserves the peripheral structures of L and L' (i.e. L and L' are the same links). By C. McA. Gordon and J. Luecke [GL], every knot in S^3 has this property. It is easy to see that a link with a non-split unknotted component does not have a unique exterior. In particular, every Milnor link does not have a unique exterior. Recently A. Kawachi [Kw3] proved by using the imitation theory [Kw1, Kw2] that the statement of the Poincaré conjecture is equivalent to the statement that a link L in S^3 having a unique exterior has Property FP.

From Theorem 1.1 and Theorem 1.2, we have a corollary which implies that the Milnor links with $\lambda \geq 3$ has Property FP' and Property FR.

Corollary 1.3 *For $\lambda \geq 3$, any finite slope surgery along M_λ does not yield both S^3 and $S^1 \times S^2$.*

Since the proof of Corollary 1.3 comes down to the proof of non-triviality of a certain knot in S^3 by the result of P. Kronheimer and T. Mrowka [KM], M_λ with $\lambda \geq 3$ has Property FP. B. Mangum and T. Stanford [MS] proved that every non-trivial Brunnian link is determined by its complement. We notice that it does not imply that every non-trivial Brunnian link has a unique exterior, but implies that if a link L in S^3 has a homeomorphic complement to a Brunnian link L' , then L is equivalent to L' or L is not a Brunnian link. Therefore Corollary 1.3 does not contribute to an alternative proof of the Poincaré conjecture. However it provides the cyclic surgery problem of links with sensitive examples.

A λ -component link in S^3 is a *Brunnian type link* if the link is an algebraically split link such that every component is an unknot and the Alexander polynomial of every proper sublink with at least two components is zero (Subsection 6.1). In Section 6, we generalize our results for the cases of Brunnian type links (Theorem 6.1, Theorem 6.7 and Corollary 6.8), and toroidal Brunnian type links (Corollary 6.16 (2)) which are Brunnian type links including essential tori in the link complements (Subsection 6.2). Moreover we characterized toroidal Brunnian links and toroidal Brunnian type links in some senses.

In Section 2, we define the Reidemeister torsions, and state their surgery formulae and properties of Alexander polynomials which are basic tools in the present paper. In Section 3, we show some key lemmas for the proofs of Theorem 1.1, Theorem 1.2, and their generalizations in Section 6 by the techniques in [Kd1, Kd2, Kd3, KMS, Tr1, Tr2] on the Reidemeister torsions. In Section 4, we prove Theorem 1.1. In Section 5, we prove Theorem 1.2 and Corollary 1.3. A necessary condition for existence of a lens surgery

is obtained from a lemma in Section 3, and non-existence of a lens surgery is shown by results due to Y. Ni [Ni] (Theorem 5.2) on fiberedness of a knot yielding a lens space, and D. Gabai [Gb1] (Theorem 5.3) on minimal genus Seifert surfaces for a plumbed link, and existence of an essential torus in the surgered manifold. Corollary 1.3 is shown by the proofs of Theorem 1.1 and Theorem 1.2, and results due to P. Kronheimer and T. Mrowka [KM] on Property P for a knot, D. Gabai [Gb1] on Property R for a knot. We remark that we cannot prove Theorem 1.2 by only using the Reidemeister torsions and the Casson-Walker invariants. This implies that an invariant deduced from the knot Floer homology due to L. Nicolaescu [Nc] cannot prove it. However the knot Floer homology itself can prove it by Ni's theorem above.

2 Reidemeister torsion

We define the Reidemeister torsions, and state their surgery formulae and properties of Alexander polynomials which are basic tools in the present paper.

2.1 Surgery formulae

Let X be a finite CW complex with $H = H_1(X)$, R an integral domain, and $\psi : \mathbb{Z}[H] \rightarrow Q(R)$ a ring homomorphism where $Q(R)$ is the quotient field of R . Then $\tau^\psi(X) \in Q(R)$ is the *Reidemeister torsion* of X related with ψ which is determined up to multiplication of $\pm\psi(h)$ ($h \in H$) (cf. [Tr1, Tr2]). We do not give a precise definition here. For A and $B \in Q(R)$, we denote $A \doteq B$ if $A = \pm\psi(h)B$ for some $h \in H$. If ψ is the identity map, then we denote $\tau(X) := \tau^\psi(X)$ for simplicity.

The Reidemeister torsion of X is defined from a cell chain complex $\mathbf{C}_*$ of the maximal abelian covering of X over a ring $\mathbb{Z}[H]$. We can take a basis $\mathbf{e}$ of $\mathbf{C}_*$ as an ordered oriented lifts of cells of X . Then the value of $\tau^\psi(X)$ is determined as a unique element in $Q(R)$. A base change with the determinant 1 preserves the value of $\tau^\psi(X)$, and a difference of two bases is described by an element of $\pm H_1(X)$. We define the basis $\mathbf{e}$ of $\mathbf{C}_*$ up to base changes with the determinant 1 a *combinatorial Euler structure* of X . The concept is equivalent to a *spin^c-structure* if X is a compact oriented 3-manifold. A great advantage of the concept is to unify separated values $\{\tau^\psi(X)\}$ to an element of $Q(\mathbb{Z}[H])$ via the Chinese Remainder Theorem (Suntzu Theorem), and a troublesome ambiguity vanishes. The unified value is called the *Reidemeister-Turaev torsion* or the *Reidemeister torsion with a combinatorial Euler structure* or the *maximal abelian torsion* (cf. [Tr2]).

Proposition 2.1 (1) We set $H_1(S^1) = \langle t \rangle \cong \mathbb{Z}$. Then we have $\tau(S^1) \doteq (t - 1)^{-1}$.

(2) We have $\tau(S^1 \times S^1) \doteq 1$.

(3) Let $L = K_1 \cup \cdots \cup K_\lambda$ be a λ -component link in an integral homology 3-sphere, and E_L the complement of L . We set $H_1(E_L) = \langle t_1, \dots, t_\lambda \rangle \cong \mathbb{Z}^\lambda$. Then we have

$$\tau(E_L) \doteq \begin{cases} \frac{\Delta_L(t_1)}{t_1 - 1} & (\lambda = 1), \\ \Delta_L(t_1, \dots, t_\lambda) & (\lambda \geq 2). \end{cases}$$

The following is coming from the excision property of the homology groups.

Lemma 2.2 (excision) *Let M_1 and M_2 be connected compact 3-manifolds with non-empty tori boundaries, and $M = M_1 \cup M_2$ a union of M_1 and M_2 such that $T = M_1 \cap M_2 = \partial M_1 \cap \partial M_2$ is a torus. Let $i_1 : M_1 \rightarrow M$, $i_2 : M_2 \rightarrow M$, $j_1 : T \rightarrow M_1$ and $j_2 : T \rightarrow M_2$ be natural inclusions. Let F be a field, $\psi : \mathbb{Z}[H_1(M)] \rightarrow F$ a ring homomorphism, and $\psi_1 = \psi \circ (i_1)_*$, $\psi_2 = \psi \circ (i_2)_*$ and $\rho = \psi \circ (i_1)_* \circ (j_1)_* = \psi \circ (i_2)_* \circ (j_2)_*$ the induced homomorphisms. Suppose that ρ is not a trivial map. Then we have*

$$\tau^\psi(M) \doteq \tau^{\psi_1}(M_1) \cdot \tau^{\psi_2}(M_2).$$

Let H be a finite cyclic group with order $p \geq 2$, and t a generator. For a divisor $d \geq 2$ of p , let ζ_d be a primitive d -th root of unity, $\psi_d : \mathbb{Z}[H] \rightarrow \mathbb{Q}(\zeta_d)$ a ring homomorphism such that $\psi_d(t) = \zeta_d$. From Lemma 2.2, we have the following surgery formulae which are suitable for studying the results of Dehn surgeries along links in an integral homology 3-sphere.

Lemma 2.3 (surgery formula)

(1) *Let $L = K_1 \cup \cdots \cup K_\lambda$ be a λ -component link with $\lambda \geq 2$ in an integral homology 3-sphere, ℓ'_i ($i = 1, \dots, \lambda$) a core of attaching solid torus, and Y the result of Dehn surgery along L with $\sharp(H_1(Y)) \geq 2$. Let F be a field, and $\psi : \mathbb{Z}[H_1(Y)] \rightarrow F$ a ring homomorphism. If $\psi([\ell'_i]) \neq 1$ for every $i = 1, \dots, \lambda$, then we have*

$$\tau^\psi(Y) \doteq \Delta_L(\psi([m_1]), \dots, \psi([m_\lambda])) \cdot \prod_{i=1}^{\lambda} (\psi([\ell'_i]) - 1)^{-1} \in F.$$

(2) *Let K be a knot in an integral homology 3-sphere, $Y = (K; p/q)$ ($p \geq 2$) the result of p/q -surgery along K , and t a generator of $H_1(Y) \cong \mathbb{Z}/p\mathbb{Z}$ corresponding to a meridian of K . Then we have*

$$\tau^{\psi_d}(Y) \doteq \Delta_K(\zeta_d)(\zeta_d - 1)^{-1}(\zeta_d^{\bar{q}} - 1)^{-1}$$

where $q\bar{q} \equiv 1 \pmod{p}$.

For example, we have

$$\tau^{\psi_d}(L(p, q)) \doteq (\zeta_d - 1)^{-1}(\zeta_d^{\bar{q}} - 1)^{-1}. \quad (2.1)$$

The Torres formula for the Alexander polynomial is a kind of surgery formulae. Let $L = K_1 \cup \cdots \cup K_\lambda \cup K_{\lambda+1}$ ($\lambda \geq 1$) be an oriented $(\lambda+1)$ -component link in a homology 3-sphere, $L' = K_1 \cup \cdots \cup K_\lambda$ a λ -component sublink, and $k_i = \text{lk}(K_i, K_{\lambda+1})$ ($i = 1, \dots, \lambda$).

Lemma 2.4 (Torres formula [To])

$$\Delta_L(t_1, \dots, t_\lambda, 1) \doteq \begin{cases} \frac{t_1^{k_1} - 1}{t_1 - 1} \Delta_{L'}(t_1) & (\lambda = 1), \\ (t_1^{k_1} \dots t_\lambda^{k_\lambda} - 1) \Delta_{L'}(t_1, \dots, t_\lambda) & (\lambda \geq 2). \end{cases}$$

Lemma 2.5 (duality [Mi3, Tr1]) *Let $L = K_1 \cup \dots \cup K_\lambda$ be a λ -compnent link in a homology 3-sphere. Then we have*

$$\Delta_L(t_1) = t_1^e \Delta_L(t_1^{-1}) \quad (\lambda = 1)$$

for some even integer e , and

$$\Delta_L(t_1, \dots, t_\lambda) = (-1)^{\lambda} t_1^{e_1} \dots t_\lambda^{e_\lambda} \Delta_L(t_1^{-1}, \dots, t_\lambda^{-1}) \quad (\lambda \geq 2)$$

where $e_i \equiv 1 + \sum_{j \neq i} \text{lk}(K_i, K_j) \pmod{2}$ ($i = 1, \dots, \lambda$).

2.2 d -norm

About algebraic fields, we refer the reader to L. C. Washington [Was] for example. For an element x in the d -th cyclotomic field $\mathbb{Q}(\zeta_d)$, the d -norm of x is defined as

$$N_d(x) = \prod_{\sigma \in \text{Gal}(\mathbb{Q}(\zeta_d)/\mathbb{Q})} \sigma(x),$$

where $\text{Gal}(\mathbb{Q}(\zeta_d)/\mathbb{Q})$ is the Galois group related with a Galois extension $\mathbb{Q}(\zeta_d)$ over $\mathbb{Q}$. The order of $\text{Gal}(\mathbb{Q}(\zeta_d)/\mathbb{Q})$ is the Euler function $\varphi : \mathbb{N} \rightarrow \mathbb{N}$ at d . In particular, $\varphi(1) = \varphi(2) = 1$, and $\varphi(d)$ is even for $d \geq 3$. The following is well-known.

Proposition 2.6

- (1) *If $x \in \mathbb{Q}(\zeta_d)$, then $N_d(x) \in \mathbb{Q}$. The map $N_d : \mathbb{Q}(\zeta_d) \setminus \{0\} \rightarrow \mathbb{Q} \setminus \{0\}$ is a group homomorphism.*
- (2) *If $x \in \mathbb{Z}[\zeta_d]$, then $N_d(x) \in \mathbb{Z}$.*

By easy calculations, we have the following.

Lemma 2.7

- (1) $N_d(\pm \zeta_d) = \begin{cases} \pm 1 & (d = 2), \\ 1 & (d \geq 3). \end{cases}$
- (2) $N_d(1 - \zeta_d) = \begin{cases} \ell & (d \text{ is a power of a prime } \ell \geq 2), \\ 1 & (\text{otherwise}). \end{cases}$

3 Key Lemmas

We show some key lemmas for the proofs of Theorem 1.1, Theorem 1.2, and their generalizations in Section 6 by the techniques in [Kd1, Kd2, Kd3, KMS, Tr1, Tr2] on the Reidemeister torsions.

3.1 Alexander polynomial of algebraically split links

Let $L = K_1 \cup \cdots \cup K_\lambda$ be an oriented λ -component algebraically split link with $\lambda \geq 2$. By Lemma 2.4, we have

$$\Delta_L(t_1, \dots, t_\lambda) = (t_1 - 1) \cdots (t_\lambda - 1) f(t_1, \dots, t_\lambda) \quad (3.1)$$

where $f(t_1, \dots, t_\lambda) \in \mathbb{Z}[t_1^{\pm 1}, \dots, t_\lambda^{\pm 1}]$. By Lemma 2.5, we may assume that

$$f(t_1, \dots, t_\lambda) = f(t_1^{-1}, \dots, t_\lambda^{-1}). \quad (3.2)$$

We add one component K to L such that $L_i = K_i \cup K$ ($i = 1, \dots, \lambda$) is the connected sum of K_i and the Hopf link, where the linking number of K_i and K is 1. Then we have

$$\Delta_{L_i}(t_i, t) \doteq \Delta_{K_i}(t_i). \quad (3.3)$$

We set $\bar{L} = L \cup K$. By Lemma 2.4 and (3.1), we have

$$\begin{aligned} \Delta_{\bar{L}}(t_1, \dots, t_\lambda, t) &= (t_1 \cdots t_\lambda - 1)(t_1 - 1) \cdots (t_\lambda - 1) f(t_1, \dots, t_\lambda) \\ &\quad + (t - 1) g(t_1, \dots, t_\lambda, t) \end{aligned} \quad (3.4)$$

where $g(t_1, \dots, t_\lambda, t) \in \mathbb{Z}[t_1^{\pm 1}, \dots, t_\lambda^{\pm 1}, t^{\pm 1}]$, and we may replace $f(t_1, \dots, t_\lambda)$ and $g(t_1, \dots, t_\lambda, t)$ with $-f(t_1, \dots, t_\lambda)$ and $-g(t_1, \dots, t_\lambda, t)$. By Lemma 2.5, we may assume that

$$\Delta_{\bar{L}}(t_1, \dots, t_\lambda, t) = (-1)^{\lambda+1} t_1^2 \cdots t_\lambda^2 t^a \Delta_{\bar{L}}(t_1^{-1}, \dots, t_\lambda^{-1}, t^{-1}) \quad (3.5)$$

where $a \equiv \lambda + 1 \pmod{2}$. We set $I_\lambda = \{1, 2, \dots, \lambda\}$. For $I = \{i_1, i_2, \dots, i_\mu\} \subset I_\lambda$, we set $L_I = K_{i_1} \cup \cdots \cup K_{i_\mu}$, $\bar{L}_I = L_I \cup K$, and $g_I \in \mathbb{Z}[t_{i_1}^{\pm 1}, \dots, t_{i_\mu}^{\pm 1}, t^{\pm 1}]$ is obtained by substituting $t_j = 1$ for all $j \in I_\lambda \setminus I$ to $g(t_1, \dots, t_\lambda, t)$.

By (3.3) and (3.5), if $I = \{i\}$ ($s = 1$), then we may take

$$\Delta_{\bar{L}_I}(t_i, t) = \Delta_{K_i}(t_i) \quad (3.6)$$

where $\Delta_{K_i}(t_i) = t_i^2 \Delta_{K_i}(t_i^{-1})$ and $\Delta_{K_i}(1) = 1$. If $2 \leq s \leq \lambda$ ($\lambda \geq 2$), then we may take

$$\begin{aligned} \Delta_{\bar{L}_I}(t_{i_1}, \dots, t_{i_s}, t) &= \left(\prod_{i \in I} t_i - 1 \right) \prod_{i \in I} (t_i - 1) f_I(t_{i_1}, \dots, t_{i_s}) \\ &\quad + (t - 1) g'_I(t_{i_1}, \dots, t_{i_s}, t) \end{aligned} \quad (3.7)$$

where

$$\Delta_{L_I}(t_{i_1}, \dots, t_{i_s}) = \prod_{i \in I} (t_i - 1) f_I(t_{i_1}, \dots, t_{i_s}),$$

$$f_I(t_{i_1}, \dots, t_{i_s}) = f_I(t_{i_1}^{-1}, \dots, t_{i_s}^{-1}) \in \mathbb{Z}[t_{i_1}^{\pm 1}, \dots, t_{i_\mu}^{\pm 1}]$$

and

$$g'_I(t_{i_1}, \dots, t_{i_s}, t) \in \mathbb{Z}[t_{i_1}^{\pm 1}, \dots, t_{i_\mu}^{\pm 1}, t^{\pm 1}].$$

We set $f_I = f_I(t_{i_1}, \dots, t_{i_s})$ and $g'_I = g'_I(t_{i_1}, \dots, t_{i_s}, t)$. We remark that $f_{I_\lambda} = f = f(t_1, \dots, t_\lambda)$ and $g'_{I_\lambda} = g = g(t_1, \dots, t_\lambda, t)$.

Lemma 3.1 *Under the situation above, for $1 \leq s \leq \lambda - 1$, we have*

$$g_I = (t - 1)^{\lambda-s-1} \Delta_{L_I}(t_{i_1}, \dots, t_{i_s}, t).$$

Proof. By applying Lemma 2.4 repeatedly, we have the result. ■

We expand the $g(t_1, \dots, t_\lambda, t)$ -part in (3.4) with normalization as follows:

Lemma 3.2 *For $\lambda \geq 2$, we have*

$$\begin{aligned} g(t_1, \dots, t_\lambda, t) &= (t - 1)^{\lambda-2} \prod_{i=1}^{\lambda} \Delta_{K_i}(t_i) \\ &+ \sum_{\substack{I \subset I_\lambda \\ 2 \leq s=|I| \leq \lambda-1}} \prod_{i \in I} (t_i - 1) (t - 1)^{\lambda-s-1} \\ &\cdot \left\{ \left(\prod_{i \in I} t_i - 1 \right) f_I + (t - 1) h_I \right\} \\ &+ \prod_{i=1}^{\lambda} (t_i - 1) h \end{aligned}$$

where $|I|$ is the number of elements of I , and $f_I \in \mathbb{Z}[t_{i_1}^{\pm 1}, \dots, t_{i_\mu}^{\pm 1}]$, $h_I \in \mathbb{Z}[t_{i_1}^{\pm 1}, \dots, t_{i_\mu}^{\pm 1}, t^{\pm 1}]$ and $h \in \mathbb{Z}[t_1^{\pm 1}, \dots, t_\lambda^{\pm 1}, t^{\pm 1}]$.

For the proof of Lemma 3.2, we refer the reader to [Kd3, Lemma 4.2].

Corollary 3.3 $g(1, \dots, 1, t) = (t - 1)^{\lambda-2}$.

Remark 3.4 By Corollary 3.3, the normalization of $g(t_1, \dots, t_\lambda, t)$ is uniquely determined, and the normalization of $f(t_1, \dots, t_\lambda)$ is also uniquely determined.

3.2 Key Lemmas

Let L and $\bar{L}$ be the same links as in Subsection 3.1.

We set $Y = (L; p_1/q_1, \dots, p_\lambda/q_\lambda)$ and $\bar{Y} = (\bar{L}; p_1/q_1, \dots, p_\lambda/q_\lambda, \emptyset)$ with $q_i \neq 0$ ($i = 1, \dots, \lambda$). We calculate a presentation matrix of the first homologies $H_1(Y)$ and $H_1(\bar{Y})$. Let m_i and l_i be a meridian and a longitude of K_i ($i = 1, \dots, \lambda$), and m and l a meridian and a longitude of K . Let T_i be an attaching solid torus of K_i , T an attaching solid torus of K , m'_i and l'_i a meridian and a core of T_i , and m' and l' a meridian and a core of T . We denote the representing element of a loop γ in the first homology by $[\gamma]$. Then we have

$$\begin{aligned} H_1(E_L) &= \langle [m_1], \dots, [m_\lambda] \rangle \cong \mathbb{Z}^\lambda \\ H_1(E_{\bar{L}}) &= \langle [m_1], \dots, [m_\lambda], [m] \rangle \cong \mathbb{Z}^{\lambda+1} \end{aligned}$$

and

$$\begin{aligned} H_1(Y) &= \langle [m_1], [l_1], \dots, [m_\lambda], [l_\lambda] \mid [m_i]^{p_i} [l_i]^{q_i} = 1, [l_i] = 1 \ (i = 1, \dots, \lambda) \rangle \\ &\cong \langle [m_1], \dots, [m_\lambda] \mid [m_i]^{p_i} = 1 \ (i = 1, \dots, \lambda) \rangle \\ &\cong \bigoplus_{i=1}^{\lambda} \mathbb{Z}/p_i \mathbb{Z}. \end{aligned}$$

Hence $H_1(Y) \cong \mathbb{Z}/p\mathbb{Z}$ if and only if $p = |p_1 \cdots p_\lambda|$ and $\gcd(p_i, p_j) = 1$ ($1 \leq i \neq j \leq \lambda$). From now on, we assume $p \geq 2$.

$$\begin{aligned} H_1(\bar{Y}) &= \left\langle \begin{array}{l} [m_1], [l_1], \dots, [m_\lambda], [l_\lambda], \\ [m'_1], [l'_1], \dots, [m'_\lambda], [l'_\lambda], \\ [m], [l] \end{array} \mid \begin{array}{l} [m'_i] = [m_i]^{p_i} [l_i]^{q_i} = 1, \\ [l'_i] = [m_i]^{r_i} [l_i]^{s_i} \ (p_i s_i - q_i r_i = -1), \\ [l_i] = [m] \ (i = 1, \dots, \lambda), \\ [l] = [m_1] \cdots [m_\lambda] \end{array} \right\rangle \\ &\cong \langle [m_1], \dots, [m_\lambda], [m] \mid [m_i]^{p_i} [m]^{q_i} = 1 \ (i = 1, \dots, \lambda) \rangle \\ &\cong \mathbb{Z}. \end{aligned} \tag{3.8}$$

We set $t_i = [m_i]$, $t = [m]$ in $H_1(Y)$ and $H_1(\bar{Y})$, and a generator of $H_1(\bar{Y})$ as T . Then a group homomorphism $\rho : H_1(\bar{Y}) \rightarrow \langle T \rangle \cong \mathbb{Z}$ defined by

$$\rho(t_i) = T^{\frac{q_i p}{p_i}}, \quad \rho(t) = T^{-p} \tag{3.9}$$

is a well-defined isomorphism. Then we have

$$\rho([l'_i]) = T^{\frac{p}{p_i}}, \quad \rho([l]) = \rho(t_1 \cdots t_\lambda) = T^{\frac{q_1 p}{p_1} + \cdots + \frac{q_\lambda p}{p_\lambda}}. \tag{3.10}$$

We note that the result of ∞ -surgery along K from $\bar{Y}$ is Y , and since it corresponds to $T^{-p} = 1$ in (3.9),

$$H_1(Y) \cong \langle T \mid T^p = 1 \rangle \cong \mathbb{Z}/p\mathbb{Z}$$

is recovered.

For fixed k , we set $f_k = f_k(t_k) = f(1, \dots, 1, t_k, 1, \dots, 1)$ which is obtained by substituting $t_j = 1$ ($j \neq k$) to $f(t_1, \dots, t_\lambda)$. Then we have the following:

Lemma 3.5 *Suppose that L is a λ -component Brunnian link with $\lambda \geq 3$. For a divisor $d \geq 2$ of p_k , we have*

$$\tau^{\psi_d}(Y) \doteq \left\{ f_k(\zeta_d) \left(\prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} q_j \right) (\zeta_d - 1)^2 + (-1)^\lambda \left(\prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} p_j \right) \zeta_d \right\} \cdot (\zeta_d - 1)^{-1} (\zeta_d^{\bar{q}_k} - 1)^{-1}$$

where $q_k \bar{q}_k \equiv 1 \pmod{p_k}$.

Proof. Since L is a Brunnian link, we have $\Delta_{K_i}(t_i) = t_i$ and $f_I = 0$ in Lemma 3.2. By Lemma 2.3 (1), Lemma 3.2, (3.9) and (3.10), we have

$$\begin{aligned} \tau(\bar{Y}) &\doteq \Delta_{\bar{L}}(\rho(t_1), \dots, \rho(t_\lambda), \rho(t)) \prod_{i=1}^{\lambda} (\rho([l'_i]) - 1)^{-1} \\ &\doteq \left(T^{\frac{q_1 p}{p_1} + \dots + \frac{q_\lambda p}{p_\lambda}} - 1 \right) \prod_{i=1}^{\lambda} \frac{T^{\frac{q_i p}{p_i}} - 1}{T^{\frac{p}{p_i}} - 1} \cdot f \left(T^{\frac{q_1 p}{p_1}}, \dots, T^{\frac{q_\lambda p}{p_\lambda}} \right) \\ &\quad + (-1)^\lambda T^{p\lambda + \frac{q_1 p}{p_1} + \dots + \frac{q_\lambda p}{p_\lambda}} \prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} \frac{T^p - 1}{T^{\frac{p}{p_j}} - 1} \cdot \left(T^{\frac{p}{p_k}} - 1 \right)^{-1} \\ &\quad + (T^{-p} - 1) \sum_{\substack{I \subset I_\lambda \\ 2 \leq s=|I| \leq \lambda-1}} \prod_{i \in I} \frac{T^{\frac{q_i p}{p_i}} - 1}{T^{\frac{p}{p_i}} - 1} \prod_{j \in I_\lambda \setminus I} \frac{T^{-p} - 1}{T^{\frac{p}{p_j}} - 1} \\ &\quad \cdot h_I \left(T^{\frac{q_{i_1} p}{p_{i_1}}}, \dots, T^{\frac{q_{i_\mu} p}{p_{i_\mu}}}, T^{-p} \right) \\ &\quad + (T^{-p} - 1) \prod_{i=1}^{\lambda} \frac{T^{\frac{q_i p}{p_i}} - 1}{T^{\frac{p}{p_i}} - 1} \cdot h \left(T^{\frac{q_1 p}{p_1}}, \dots, T^{\frac{q_\lambda p}{p_\lambda}}, T^{-p} \right). \end{aligned} \tag{3.11}$$

Since p/p_k and q_k are relatively prime to p_k , $\frac{q_k p}{p_k}$ is also relatively prime to p_k . We define a ring homomorphism $\bar{\psi}_d : \mathbb{Z}[T, T^{-1}] \rightarrow \mathbb{Q}(\zeta_d)$ by

$$\bar{\psi}_d(T) = \zeta_d^{\frac{\overline{q_k p}}{p_k}}$$

where $\frac{q_k p}{p_k} \cdot \frac{\overline{q_k p}}{p_k} \equiv 1 \pmod{p_k}$. Then we have

$$\bar{\psi}_d \left(T^{\frac{q_i p}{p_i}} \right) = \bar{\psi}_d \left(T^{\frac{p}{p_i}} \right) = 1 \quad (i \neq k), \quad \bar{\psi}_d \left(T^{\frac{q_k p}{p_k}} \right) = \zeta_d, \quad \bar{\psi}_d \left(T^{\frac{p}{p_k}} \right) = \zeta_d^{\bar{q}_k}, \quad \bar{\psi}_d(T^p) = 1,$$

$$\begin{aligned}\overline{\psi}_d \left(\frac{T^{\frac{q_i p}{p_i}} - 1}{T^{\frac{p}{p_i}} - 1} \right) &= \text{sign}(q_i) \cdot \overline{\psi}_d \left(\sum_{j=1}^{|q_i|-1} T^{\frac{j p}{p_i}} \right) = q_i \quad (i \neq k), \\ \overline{\psi}_d \left(\frac{T^p - 1}{T^{\frac{p}{p_i}} - 1} \right) &= \text{sign}(p_i) \cdot \overline{\psi}_d \left(\sum_{j=1}^{|p_i|-1} T^{\frac{j p}{p_i}} \right) = p_i \quad (i \neq k),\end{aligned}$$

and by (3.11), we have

$$\tau^{\overline{\psi}_d}(\overline{Y}) \doteq \left\{ f_k(\zeta_d) \left(\prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} q_j \right) (\zeta_d - 1)^2 + (-1)^\lambda \left(\prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} p_j \right) \zeta_d \right\} \cdot (\zeta_d^{\overline{q}_k} - 1)^{-1}. \quad (3.12)$$

By Lemma 2.2, $\psi_d([l]) = \zeta_d$, and (3.12), we have

$$\begin{aligned}\tau^{\psi_d}(Y) &\doteq \tau^{\overline{\psi}_d}(\overline{Y}) \cdot (\zeta_d - 1)^{-1} \\ &= \left\{ f_k(\zeta_d) \left(\prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} q_j \right) (\zeta_d - 1)^2 + (-1)^\lambda \left(\prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} p_j \right) \zeta_d \right\} \\ &\quad \cdot (\zeta_d - 1)^{-1} (\zeta_d^{\overline{q}_k} - 1)^{-1}.\end{aligned}$$

■

Lemma 3.6 *In the same situation as in Lemma 3.5, we suppose Y is a lens space. Then we have the following:*

- (1) *If f_k is constant with $f_k \geq 1$ and $|p_k| \geq 5$, then we have $f_k = 1$, $|q_j| = 1$ ($j \neq k$), and $\prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} p_j = \eta$, 2η or 3η where $\eta = \prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} q_j = 1$ or -1 .*

- (2) *If $f_k = 0$, then we have $|p_j| = 1$ ($j \neq k$).*

Proof. Since Y is a lens space and (2.1), we have

$$\tau^{\psi_d}(Y) \doteq (\zeta_d^a - 1)^{-1} (\zeta_d^b - 1)^{-1}$$

for some a and b , where $\gcd(a, p_k) = \gcd(b, p_k) = 1$. Then by Lemma 3.5, we have an equation

$$\left\{ f_k(\zeta_d) \left(\prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} q_j \right) (\zeta_d - 1)^2 + (-1)^\lambda \left(\prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} p_j \right) \zeta_d \right\} \cdot (\zeta_d^a - 1)(\zeta_d^b - 1) \doteq (\zeta_d - 1)(\zeta_d^c - 1) \quad (3.13)$$

where $c \equiv \pm \bar{q}_k \pmod{p_k}$.

By fixing a combinatorial Euler structure, we can lift the equation above to the equation in $\mathbb{Z}[u, u^{-1}]/(u^{|p_k|-1} + \dots + u + 1)$ as follows :

$$\left\{ f_k(u) \left(\prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} q_j \right) (u-1)^2 + (-1)^\lambda \left(\prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} p_j \right) u \right\} \cdot (u^a-1)(u^b-1) = \pm u^l (u-1)(u^c-1) \quad (3.14)$$

and it deduces the following :

$$\left\{ f_k(u) \left(\prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} q_j \right) (u-1)^2 + (-1)^\lambda \left(\prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} p_j \right) u \right\} \cdot \frac{u^a-1}{u-1} \cdot \frac{u^b-1}{u-1} = \pm u^l \cdot \frac{u^c-1}{u-1}. \quad (3.15)$$

We set the part $\{\dots\}$ in (3.14) and (3.15) as $F(u)$. If f_k is constant, then $F(u)$ is of degree 2. We set $F(u) = mu^2 + nu + m$ ($m, n \in \mathbb{Z}$). By taking the d -norm (cf. Subsection 2.2) of both sides of (3.13), we have $(m, n) \neq (0, 0)$ and $\gcd(m, n) = 1$ (The d -norm of the part $\{\dots\}$ in (3.13) is 1).

(1) Suppose that f_k is constant with $f_k \geq 1$ and $|p_k| \geq 5$.

(i) The case that p_k is even.

We can take $1 \leq a, b, c \leq |p_k|/2 - 1$ are odd, the degree of the lefthand side of (3.15) is $\leq |p_k| - 2$, and the degree of the righthand side of (3.15) is $\leq |p_k|/2 - 2$. Since both lefthand side and righthand side of (3.15) are symmetric polynomials, they are identical as elements in $\mathbb{Z}[u, u^{-1}]$ up to multiplications of $\pm u^l$. Hence $F(u)$ is a divisor of $(u^c - 1)/(u - 1)$, and $F(u)$ is a cyclotomic polynomial of degree 2

$$\pm(u^2 + u + 1), \quad \pm(u^2 + 1) \text{ or } \pm(u^2 - u + 1).$$

Therefore we have the result.

(ii) The case that p_k is odd.

We can take $1 \leq a, b, c \leq |p_k| - 1$ satisfying $2 \leq a + b \leq |p_k| - 1$ and $a + b \equiv c + 1 \pmod{2}$. If $a + b \leq |p_k| - 2$, then by the same argument as (i), we have the result. Suppose that $a + b = |p_k| - 1$ (i.e. $a = b = (|p_k| - 1)/2$ and c is odd with $1 \leq c \leq |p_k| - 2$).

We calculate the lefthand side of (3.14), and deform it modulo $(u^{|p_k|-1} + \dots + u + 1)$.

$$\begin{aligned}
(mu^2 + nu + m) \left(u^{\frac{|p_k|-1}{2}} - 1 \right)^2 &= (mu^2 + nu + m) \left(u^{|p_k|-1} - 2u^{\frac{|p_k|-1}{2}} + 1 \right) \\
&= mu^{|p_k|+1} + nu^{|p_k|} + mu^{|p_k|-1} - 2mu^{\frac{|p_k|+3}{2}} - 2nu^{\frac{|p_k|+1}{2}} - 2mu^{\frac{|p_k|-1}{2}} \\
&\quad + mu^2 + nu + m \\
&= u^{\frac{|p_k|+1}{2}} \left\{ -2n - 2m(u + u^{-1}) + m \left(u^{\frac{|p_k|-3}{2}} + u^{-\frac{|p_k|-3}{2}} \right) + (m+n) \left(u^{\frac{|p_k|-1}{2}} + u^{-\frac{|p_k|-1}{2}} \right) \right\} \\
&= u^{\frac{|p_k|+1}{2}} \{ (-m-3n) + (-3m-n)(u + u^{-1}) - (m+n)(u^2 + u^{-2}) - \dots \\
&\quad - (m+n) \left(u^{\frac{|p_k|-5}{2}} + u^{-\frac{|p_k|-5}{2}} \right) - n \left(u^{\frac{|p_k|-3}{2}} + u^{-\frac{|p_k|-3}{2}} \right) \}.
\end{aligned} \tag{3.16}$$

(a) The case $3 \leq c \leq |p_k| - 4$ ($|p_k| \geq 7$).

The lefthand side of (3.14) is $\pm u^l(u^{c+1} - u^c - u + 1)$. By (3.16), we have $-m-3n = 0$, and $(m, n) = (3, -1)$ or $(-3, 1)$. However $-3m - n = -8$ or 8 . Hence we have no root for (m, n) .

(b) The case $c = 1$.

The lefthand side of (3.14) is $\pm u^l(u^2 - 2u + 1)$. By (3.16), we have $-m-3n = \mp 2$. If $|p_k| \geq 7$, then we have $-3m - n = \pm 1$ and $n = 0$. Hence we have no root for (m, n) . If $|p_k| = 5$, then we have $-3m - 2n = \pm 1$. Hence we have $(m, n) = (-1, 1)$ or $(1, -1)$.

(c) The case $c = |p_k| - 2$.

The lefthand side of (3.14) is

$$\pm u^l(u^{|p_k|-1} - u^{|p_k|-2} - u + 1) = \mp u^{l+1}(2u^{|p_k|-3} + u^{|p_k|-4} + \dots + u + 2).$$

By (3.16), we have $-m-3n = \pm 1$. If $|p_k| \geq 7$, then we have $-n = \pm 2$ and $-m-n = \pm 1$. Hence we have no root for (m, n) . If $|p_k| = 5$, then we have $-3m - 2n = \pm 2$. Hence we have no root for (m, n) .

Therefore we have the result.

(2) Suppose that $f_k = 0$.

The equation (3.13) is simplified as

$$\left(\prod_{\substack{j \neq k \\ 1 \leq j \leq \lambda}} p_j \right) (\zeta_d^a - 1)(\zeta_d^b - 1) \doteq (\zeta_d - 1)(\zeta_d^c - 1).$$

By taking the d -norm of both sides, we have the result. ■

By (1.1), the Alexander polynomial of M_3 satisfies $f = f_k = 1$ corresponding to Lemma 3.6 (1), and that of M_λ ($\lambda \geq 4$) satisfies $f = f_k = 0$ corresponding to Lemma 3.6 (2).




Figure 2: n -twisted Whitehead link W_n

Lemma 3.7 [KMS, Theorem 1.2] *Let W_n ($n \neq 0$) be the n -twisted Whitehead link as in Figure 2 where the rectangle implies $(-n)$ -full twists, and $Y = (W_n; p_1/q_1, p_2/q_2)$ the result of a finite slope surgery along W_n .*

- (a) *If $|n| \geq 2$, then Y is not a lens space.*
- (b) [MP] *Suppose that $n = \varepsilon = 1$ or -1 . Then Y is a lens space if and only if one of the following holds:*

- (1) $p_1/q_1 = \varepsilon$ and $|\varepsilon p_2 - 6q_2| = 1$, or
- (2) $p_1/q_1 = 2\varepsilon$ and $|\varepsilon p_2 - 4q_2| = 1$, or
- (3) $p_1/q_1 = 3\varepsilon$ and $|\varepsilon p_2 - 3q_2| = 1$,

and the cases that indices of (p_i, q_i) ($i = 1, 2, 3$) are permuted. Moreover, if (1), then $Y = L(p_2, 4\varepsilon q_2)$, if (2), then $Y = L(2p_2, \varepsilon(8q_2 - p_2))$, and if (3), then $Y = L(3p_2, \varepsilon(3q_2 - 2p_2))$.

4 Proof of Theorem 1.1

Since the components of M_3 are interchangeable one another, we may suppose $1 \leq |p_1| \leq |p_2| \leq |p_3|$. By Lemma 3.6 (1), we have $|p_1| = 1$.

- (i) The case $|p_3| \geq 5$.

We set $p_1/q_1 = \varepsilon$. Since $(M_3; \varepsilon, p_2/q_2, p_3/q_3) = (W_\varepsilon; p_2/q_2, p_3/q_3)$ and Lemma 3.7 (b), we have the result.

- (ii) The case $|p_3| \leq 4$.

We set $p_1/q_1 = 1/n$ ($n \neq 0$). Since $(M_3; 1/n, p_2/q_2, p_3/q_3) = (W_n; p_2/q_2, p_3/q_3)$ and Lemma 3.7 (a), we have $n = \pm 1$. By the similar argument as (i), we have the result. ■

5 Proofs of Theorem 1.2 and Corollary 1.3

We suppose that L and $\bar{L}$ are the same links as in Section 3, $\lambda \geq 4$, and $|p_i| \geq 2$ for fixed i . By Lemma 3.6 (2), we may assume $p_j = 1$ ($j \neq i$). Let $\widehat{K}_i$ be the resulting knot by $1/q_j$ -surgery along K_j ($j \neq i$) of L , and $\bar{L}_i$ the resulting 2-component link by $1/q_j$ -surgery along K_j ($j \neq i$) of $\bar{L}$. We set $Y_i = E_{\widehat{K}_i}$ and $\bar{Y}_i = E_{\bar{L}_i}$. We note that $\widehat{K}_i$ and $\bar{L}_i$ are a knot and a link in S^3 . Then we have

$$\begin{aligned}
H_1(Y_i) &= \langle [m_1], [l_1], \dots, [m_\lambda], [l_\lambda] \mid [m_j][l_j]^{q_j} = 1, [l_j] = 1 \ (j \neq i), [l_i] = 1 \rangle \\
&\cong \langle [m_1], \dots, [m_\lambda] \mid [m_j] = 1 \ (j \neq i) \rangle \\
&\cong \langle [m_i] \mid - \rangle \cong \mathbb{Z}, \\
H_1(\bar{Y}_i) &= \left\langle \begin{array}{l} [m_1], [l_1], \dots, [m_\lambda], [l_\lambda], \\ [m'_1], [l'_1], \dots, [m'_\lambda], [l'_\lambda], \\ [m], [l] \end{array} \mid \begin{array}{l} [m'_j] = [m_j][l_j]^{q_j} = 1, \\ [l'_j] = [l_j]^{-1} \ (j \neq i), \\ [l_j] = [m] \ (j = 1, \dots, \lambda), \\ [l] = [m_1] \cdots [m_\lambda] \end{array} \right\rangle \\
&\cong \langle [m_1], \dots, [m_\lambda], [m] \mid [m_j][m]^{q_j} = 1 \ (j \neq i) \rangle \\
&\cong \langle [m_i], [m] \mid - \rangle \cong \mathbb{Z}^2,
\end{aligned} \tag{5.1}$$

and

$$\begin{aligned}
\tau(\bar{Y}_i) &\doteq \Delta_{\bar{L}}(t^{-q_1}, \dots, t_i, \dots, t^{-q_\lambda}, t) \prod_{\substack{j \neq i \\ 1 \leq j \leq \lambda}} ([l'_j] - 1)^{-1} \\
&\doteq (t-1)g(t^{-q_1}, \dots, t_i, \dots, t^{-q_\lambda}, t) \prod_{\substack{j \neq i \\ 1 \leq j \leq \lambda}} (t-1)^{-1} \\
&= t_i t^{-q_1 - \dots - \hat{q}_i - \dots - q_\lambda} \\
&\quad + (t-1)(t_i-1) \sum_{\substack{i \in I \subset I_\lambda \\ 2 \leq s=|I| \leq \lambda-1}} \prod_{j \in I \setminus \{i\}} \frac{t^{-q_j} - 1}{t-1} h_I \\
&\quad + (t-1)^2 \sum_{\substack{i \notin I \subset I_\lambda \\ 2 \leq s=|I| \leq \lambda-1}} \prod_{j \in I} \frac{t^{-q_j} - 1}{t-1} h_I \\
&\quad + (t-1)(t_i-1) \prod_{\substack{j \neq i \\ 1 \leq j \leq \lambda}} \frac{t^{-q_j} - 1}{t-1} h
\end{aligned} \tag{5.2}$$

where h_I and h are the same as in Lemma 3.2.

Lemma 5.1 $\Delta_{\widehat{K}_i}(t_i) \doteq 1$.

Proof. Let $\pi : \mathbb{Z}[t_i^{\pm 1}, t^{\pm 1}] \rightarrow \mathbb{Z}[t_i, t_i^{-1}]$ be a ring homomorphism defined by $\pi(t_i) = t_i$ and $\pi(t) = 1$. Then by (5.1) and (5.2), we have

$$\tau(Y_i) \doteq \tau^\pi(\overline{Y}_i)(\pi([l]) - 1)^{-1} = \frac{t_i}{t_i - 1}.$$

By Proposition 2.1 (3), we have the result. ■

Theorem 5.2 [Ni, Corollary 1.3] *If a knot K in S^3 yields a lens space, then K is fibered.*

By this theorem, existence of lens surgery along $\widehat{K}_i$ is equivalent to triviality of $\widehat{K}_i$.

Theorem 5.3 [Gb1] *Let F be a Seifert surface of a link L which is a Murasugi sum of two surfaces F_1 and F_2 . Then F is a minimal genus Seifert surface of L if and only if both F_1 and F_2 are minimal genus Seifert surfaces of links.*

Proof of Theorem 1.2.

(1) The case $\lambda = 4$.

By the symmetry of M_4 as in Section 1, it is sufficient to prove that $\widehat{K}_1$ is non-trivial. Let F be a genus 1 Seifert surface of M_4 as in Figure 3. Then F is a plumbing of F_1 and F_2 in Figure 3. In Figure 3, the number in a rectangle implies the number of full twists. Since ∂F_1 is a non-trivial $(2, -2q_2)$ -torus link, F_1 is a minimal genus Seifert surface. Since $\partial F_1 = K'_1 \cup K'_2$ is a parallel of a non-trivial 2-bridge knot $C(-2q_3, 2q_4)$, F_2 is a minimal genus Seifert surface. By Theorem 5.3, F is a minimal genus Seifert surface, and hence $\widehat{K}_1$ is a non-trivial knot.

(2) The case $\lambda \geq 5$.

Let Y be the result of Dehn surgery along M_λ whose surgery coefficient of K_j ($j \neq i$) is $1/q_j$ and that of K_i is p/q . We show that Y includes an essential torus. Let T be a standard torus embedded in S^3 which divides S^3 into two solid tori Σ_1 and Σ_2 . We can take T such that Σ_1 includes $L_1 = K_1 \cup K_2$, and Σ_2 includes $L_2 = K_3 \cup \cdots \cup K_\lambda$ (Figure 4). We show that both the result of Dehn surgery along L_1 in Σ_1 , which is denoted by W_1 , and the result of Dehn surgery along L_2 in Σ_2 , which is denoted by W_2 , are not solid tori.

Suppose that W_1 is a solid torus. Then in Theorem 1.1, there exists one of three sequences such that a surgery coefficients of some component can be any rational number. Hence W_1 is not a solid torus.

Suppose that W_2 is a solid torus. Since the complement of L_2 in Σ_2 is homeomorphic to $M_{\lambda-1}$ with preserving the peripheral structures of the components except $\partial \Sigma_2$, it contradicts the assumption of induction. Hence W_2 is not a solid torus. ■

Remark 5.4 (1) In the proof of Theorem 1.2, the case $\lambda = 4$ can also be proved in a similar way as the case $\lambda \geq 5$. It will be discussed again in Subsection 6.2.




Figure 3: minimal genus Seifert surface for $\hat{K}_1$




Figure 4: essential torus T , and solid tori Σ_1 and Σ_2

- (2) We cannot prove Theorem 1.2 by only using the Reidemeister torsions and the Casson-Walker invariants [Les, Wa]. Hence an invariant induced from the knot Floer homology due to L. Nicolaescu [Nc] also cannot prove it. However the knot Floer homology itself can prove it by Ni's theorem above.

Proof of Corollary 1.3. Suppose that $H_1(Y) \cong \{0\}$ (resp. $H_1(Y) \cong \mathbb{Z}$). There exists $i \in \{1, 2, \dots, \lambda\}$ such that a surgery coefficient of K_j with $j \neq i$ is $1/q_j$ and that of K_i is $1/q_i$ (resp. 0). Then since $\widehat{K}_i$ is a 2-bridge knot of genus 1 for $\lambda = 3$ and $\widehat{K}_i$ is non-trivial by the proof of Theorem 1.2 for $\lambda \geq 4$, Y is not S^3 (resp. not $S^1 \times S^2$) by the affirmative answer of Property P conjecture (resp. Property R conjecture) due to P. Kronheimer and T. Mrowka [KM] (resp. D. Gabai [Gb2]). ■

6 Generalization to Brunnian type links and toroidal Brunnian links

We generalize Theorem 1.1, Theorem 1.2 and Corollary 1.3 for the cases of *Brunnian type links* and toroidal Brunnian type links which are Brunnian links including essential tori in the link complement. Moreover we characterized toroidal Brunnian links and toroidal Brunnian type links.

6.1 Generalization to Brunnian type links

A λ -component link $L = K_1 \cup \dots \cup K_\lambda$ in S^3 with $\lambda \geq 2$ is a *Brunnian type link* if L is an algebraically split link such that every component is an unknot and the Alexander polynomial of every proper sublink with at least two components is zero. Let $\overline{L} = L \cup K$ be the same link as in Subsection 3.1. Then we have the Alexander polynomial of $\widehat{K}_i$ in Section 5 by a similar way as (5.2) and the proof of Lemma 5.1 :

$$\Delta_{\widehat{K}_i}(t_i) \doteq t_i + (-1)^{\lambda-1} \left(\prod_{\substack{j \neq i \\ 1 \leq j \leq \lambda}} q_j \right) f_i(t_i)(t_i - 1)^2. \quad (6.1)$$

We note that by Remark 3.4, the sign of $f_i(t_i)$ is uniquely determined. By (6.1) and the affirmative answer of Property P conjecture (resp. Property R conjecture) due to P. Kronheimer and T. Mrowka [KM] (resp. D. Gabai [Gb2]), we have the following.

Theorem 6.1 *We suppose the situation above (i.e. L is a λ -component Brunnian type link).*

- (1) *If the result of a finite slope surgery along L is S^3 , then $f_i = 0$ for all $i \in \{1, 2, \dots, \lambda\}$.*
- (2) *If the result of a finite slope surgery along L is $S^1 \times S^2$, then $f_i = 0$ for some $i \in \{1, 2, \dots, \lambda\}$.*

Let $L = K_1 \cup \dots \cup K_\lambda$ be a λ -component link in S^3 and $W_1 = K'_1 \cup K'_2$ the Whitehead link (cf. Figure 2). Since K'_2 is an unknot, the complement of K'_2 is a solid torus Σ and we may regard W_1 as a knot K'_1 in Σ . If we attach Σ to $\partial N(K_i)$ by identifying the preferred longitude of K'_2 with the meridian of K_i and the meridian of K'_2 with the preferred longitude of K_i , then we obtain a new λ -component link L' in S^3 such that K_i is replaced with K'_1 . We call L' a *Whitehead double* of L on K_i .

Example 6.2 Let $L = K_1 \cup K_2 \cup K_3$ be a 3-component link obtained by taking a Whitehead double on one component of the Whitehead link and taking 2-parallel of the other component of the Whitehead link. We suppose that K_1 is the Whitehead double part and $K_2 \cup K_3$ is the 2-parallel part. Then we can easily see that L is a Brunnian type link, but is not a Brunnian link satisfying $\Delta_L(t_1, t_2, t_3) = 0$. Since $(L; \emptyset, 1/q, -1/q)$ is an unknot for every integer q , L has not Property FP' and Property FR .




Figure 5: Brunnian type link $L = K_1 \cup K_2 \cup K_3$

We do not have any counterexamples for the following problems.

Question 6.3 (1) Does any non-split λ -component Brunnian type link L without Property FP' or Property FR satisfy $\Delta_L(t_1, \dots, t_\lambda) = 0$?

(2) Does any non-split anannular Brunnian type link have Property FP' or Property FR ?

Corollary 1.3 is a partial answer for Question 6.3 (2).

About lens surgeries along a knot, P. Ozsváth and Z. Szabó [OS] obtained the following by the knot Floer homology.

Theorem 6.4 (Ozsváth-Szabó [OS, Corollary 1.3]) Let K be a knot in S^3 . If K yields a lens space, then the Alexander polynomial of K is of the following form :

$$\Delta_K(t) \doteq (-1)^m + \sum_{k=1}^m (-1)^{k-1} (t^{n_k} + t^{-n_k}) \quad (n_1 > n_2 > \dots > n_m \geq 1)$$

where n_k ($k = 1, \dots, m$) is an integer.

The author and Y. Yamada [KY] obtained the following by the Reidemeister torsion.

Theorem 6.5 [KY] *Let K be a knot in an integral homology 3-sphere. If $(K; p/q)$ ($p \geq 2$) is a lens space, then the Alexander polynomial of K is of the following form :*

$$\Delta_K(t) \doteq \frac{(t^{rs} - 1)(t - 1)}{(t^r - 1)(t^s - 1)} \quad \left(\text{mod } \frac{t^p - 1}{t - 1} \right)$$

where $\gcd(r, s) = 1$ and $qrs \equiv \pm 1 \pmod{p}$.

By combining Theorem 6.4 and Theorem 6.5, M. Tange [Ta] obtained the following. M. Hedden and L. Watson [HW] obtained the same result independently.

Theorem 6.6 (Tange [Ta, Theorem 1.3], Hedden and Watson [HW, Corollary 9]) *Under the same situation as Theorem 6.4, we have $n_1 - n_2 = 1$. That is, the Alexander polynomial of K is of the following form :*

$$\Delta_K(t) \doteq t^{n_1} - t^{n_1-1} + \dots - t^{-n_1+1} + t^{-n_1}.$$

Theorem 6.6 implies that the trace, which is the sum of the roots, of $\Delta_K(t)$ is 1.

Theorem 6.7 *We suppose the same situation in Theorem 6.1. If the result of a finite slope surgery along L is a lens space, then we have the following:*

- (1) *For some $i \in \{1, 2, \dots, \lambda\}$ and every $j \in \{1, 2, \dots, \lambda\} \setminus \{i\}$, we have $|q_j| = 1$.*
- (2) *For the same $i \in \{1, 2, \dots, \lambda\}$ in (1), $\Delta_{\widehat{K}_i}(t_i)$ satisfies the conditions in Theorem 6.4, Theorem 6.5 and Theorem 6.6.*

By (6.1), Theorem 6.1 and Theorem 6.7, we have :

Corollary 6.8 *Under the same situation in Theorem 6.7, if $f_i(t_i)$ is constant, then we have (i) $f_i = 0$ and $\widehat{K}_i$ is an unknot, or (ii)*

$$f_i(t_i) = (-1)^{\lambda-1} \left(\prod_{\substack{j \neq i \\ 1 \leq j \leq \lambda}} q_j \right) = 1 \quad (\text{i.e. } |q_j| = 1 \text{ for every } j \in \{1, 2, \dots, \lambda\} \setminus \{i\}),$$

$$\Delta_{\widehat{K}_i}(t_i) \doteq t_i^2 - t_i + 1,$$

and $\widehat{K}_i$ is the trefoil.

6.2 Generalization to toroidal Brunnian type links

A link L is *toroidal* if there exists an essential torus in the complement of L . A link L is *atoroidal* if L is not toroidal.

Let $L = K_1 \cup \cdots \cup K_p$ be a p -component link in S^3 , and $L' = K'_1 \cup \cdots \cup K'_q$ a q -component link in S^3 . Suppose that for some j , the j -th component K'_j of L' is an unknot. Then the complement of K'_j is a solid torus, and we denote it by Σ' . If we attach Σ' to the complement of L by identifying the preferred longitude of K'_j with the meridian of K_i and the meridian of K'_j with the preferred longitude of K_i , then we obtain a new $(p+q-2)$ -component link in S^3 . We call the link the *satellite* of L along K_i with *pattern* (L', K'_j) or L' of type (p, q) . We denote it by $S_{ij}(L, L')$ or $S(L, L')$ simply. If we set the attaching map $f : \Sigma' \rightarrow S^3$, then $S_{ij}(L, L') = (L \setminus K_i) \cup f(L' \setminus K'_j)$. If L' is the Whitehead link, then $S_{ij}(L, L')$ is the *Whitehead double* of L , if L' is the Borromean rings, then $S_{ij}(L, L')$ is the *Bing double* of L , and if L is the Hopf link, then $S_{ij}(L, L') = L'$. In particular, M_3 is the Bing double of the Hopf link, and M_λ ($\lambda \geq 4$) is the Bing double of $M_{\lambda-1}$.

A λ -component link $L = K_1 \cup \cdots \cup K_\lambda$ in S^3 is a *semi-Brunnian link* if it is a Brunnian link for $\lambda \geq 3$, every component of it is an unknot for $\lambda = 2$, and it is an arbitrary knot for $\lambda = 1$. Then we have the following.

Lemma 6.9 (1) Let $L = K_1 \cup \cdots \cup K_p$ be a p -component semi-Brunnian link, and $L' = K'_1 \cup \cdots \cup K'_q$ a q -component Brunnian link such that $p = 1$ and $q \geq 3$, or $p \geq 2$ and $q \geq 2$. Then $S_{ij}(L, L')$ is a $(p+q-2)$ -component Brunnian link in S^3 for any $i \in \{1, 2, \dots, p\}$ and $j \in \{1, 2, \dots, q\}$.

(2) In (1), $S_{ij}(L, L')$ is non-split if and only if both L and L' are non-split.

(3) In (1), if L is non-split atoroidal and is neither the unknot nor the Hopf link, and L' is non-split atoroidal, then $S_{ij}(L, L')$ is a $(p+q-2)$ -component non-split toroidal Brunnian link in S^3 for any $i \in \{1, 2, \dots, p\}$ and $j \in \{1, 2, \dots, q\}$ with a unique essential torus.

Proof. (1) Let T be an embedded solid torus in S^3 such that T separates $S_{ij}(L, L')$ into two parts $L \setminus K_i$ and $L' \setminus K'_j$, and T separates S^3 into two parts Σ and Σ' where Σ is the complement of K_i and Σ' is the complement of K'_j . We note that Σ' is a solid torus, and Σ is also a solid torus if $p \geq 2$ and is never a solid torus if $p = 1$.

Suppose that $p \geq 2$. For $r \in \{1, 2, \dots, p\} \setminus \{i\}$, we set $L_r = L \setminus K_r$. If $p = 2$, then L_r is the empty set. Then we have $S_{ij}(L, L') \setminus K_r = S_{ij}(L_r, L')$. Since L_r is a $(p-1)$ -component trivial link and $L' \setminus K'_j$ is a $(q-1)$ -component trivial link, $S_{ij}(L, L') \setminus K_r$ is a $(p+q-3)$ -component trivial link.

For $s \in \{1, 2, \dots, q\} \setminus \{j\}$, we set $L'_s = L' \setminus K'_s$. Then we have $S_{ij}(L, L') \setminus K'_s = S_{ij}(L, L'_s)$. Since L'_s is a $(q-1)$ -component trivial link and $L \setminus K_i$ is a $(p-1)$ -component trivial link, $S_{ij}(L, L') \setminus K'_s$ is a $(p+q-3)$ -component trivial link.

(2) We use the same notations as in the proof of (1). If L or L' is split, then it is easy to see that $S_{ij}(L, L')$ is also split.

Suppose that $S_{ij}(L, L')$ is split. Let F be an embedded 2-sphere in the complement of $S_{ij}(L, L')$ (i.e. $F \cap S_{ij}(L, L') = \emptyset$) which separates $S_{ij}(L, L')$ into two non-empty links. If $T \cap F = \emptyset$, then L or L' is split. We show that the case $T \cap F \neq \emptyset$ does not occur. We suppose $T \cap F \neq \emptyset$, and T intersects with F transversally. Then $T \cap F$ consists of disjoint simple loops. We take an innermost loop γ of $T \cap F$ on F and the disk B bounded by γ on F . Then B is properly embedded in Σ or Σ' . We only show the case that B is properly embedded in Σ . If γ bounds a disk B' on T , then $B \cup B'$ bounds a 3-ball W in Σ . Since L is non-split, $W \cap L = \emptyset$ and γ can be removed by local isotopy of F along W . If γ is not null-homotopic on T , then γ is a preferred longitude of K_i . Since L is non-split, $B \cap L = B \cap (L \setminus K_i) \neq \emptyset$. It contradicts the assumption that $F \cap S_{ij}(L, L') = \emptyset$. Hence the case $T \cap F \neq \emptyset$ does not occur.

(3) We use the same notations as in the proof of (1). By (1) and (2), $S_{ij}(L, L')$ is a $(p + q - 2)$ -component non-split Brunnian link in S^3 for any $i \in \{1, 2, \dots, p\}$ and $j \in \{1, 2, \dots, q\}$.

Let E be the complement of $S_{ij}(L, L')$. Since L is neither the unknot nor the Hopf link, T neither bound a solid torus nor is boundary parallel in E . Therefore T is an essential torus in E .

We show that T is a unique essential torus in E . Suppose that there exists another essential torus T' in E (i.e. $T' \cap S_{ij}(L, L') = \emptyset$ and T' is not ambient isotopic to T in E). If $T \cap T' = \emptyset$, then T' is an essential torus in the complement of L or L' . By the assumption, the case does not occur.

We may suppose that $T \cap T' \neq \emptyset$, and T' intersects with T transversally. Then $T \cap T'$ consists of disjoint simple loops, and T' is divided into $T' \cap \Sigma$ and $T' \cap \Sigma'$ by T . We take a loop γ of $T \cap T'$. If γ bounds a disk B' on T' , then γ also bounds a disk B on T , or γ is a preferred longitude of K_i or K'_j . If γ bounds a disk B on T , then $B \cup B'$ bounds a 3-ball W in Σ or Σ' . Since both L and L' are non-split, we have $W \cap L = W \cap L' = \emptyset$, and γ can be removed by local isotopy of T' along W . If γ is a preferred longitude of K_i , then B' is properly embedded in Σ . Since L is non-split, we have $B' \cap L = B' \cap (L \setminus K_i) \neq \emptyset$. It contradicts the assumption $T' \cap S_{ij}(L, L') = \emptyset$. Hence this case does not occur. We can show that γ is not a preferred longitude of K'_j in a similar way. Therefore we may assume that $T \cap T'$ consists of parallel essential simple loops on T' , and the connected components of both $T' \cap \Sigma$ and $T' \cap \Sigma'$ are annuli. The set $T \cap T'$ is also the set of parallel essential simple loops on T . We take a connected component A' of $T' \cap \Sigma$ (resp. $T' \cap \Sigma'$), which is an annulus. Then $\partial A'$ bounds an annulus A on T (resp. T'), and $A \cup A'$ is a torus. By a slight isotopy, the torus $A \cup A'$ can be regarded as an embedded torus in $\text{Int}(\Sigma)$ (resp. $\text{Int}(\Sigma')$). Since Σ (resp. Σ') is atoroidal, the torus bounds a solid torus which is included by a regular neighborhood of $\partial \Sigma$ (resp. $\partial \Sigma'$). We can remove $\partial A'$ by local isotopy of T' along the

solid torus. Therefore we can isotope T' such that $T \cap T' = \emptyset$, and the case does not occur (T' cannot exist). ■

We show the converse of Lemma 6.9 (3) for the case $p \geq 2$ and $q \geq 2$.

Theorem 6.10 *For a λ -component non-split toroidal Brunnian link $\widehat{L}$ with $\lambda \geq 2$, if there exists an essential torus T in the complement of $\widehat{L}$ such that T decomposes $\widehat{L}$ into a $(p-1)$ -component sublink and a $(q-1)$ -component sublink with $p \geq 2$ and $q \geq 2$, then there exist a p -component non-split semi-Brunnian link L and a q -component non-split Brunnian link L' such that $\widehat{L} = S(L, L')$ with $\lambda = p + q - 2$.*

Proof. Let E be the complement of $\widehat{L}$, and T an essential torus in E . Then T decomposes S^3 into two parts Σ and Σ' , and one of Σ and Σ' is a solid torus. We suppose that Σ' is a solid torus, and that Σ includes p' components of $\widehat{L}$ and Σ' includes q' components of $\widehat{L}$. Then $\lambda = p' + q'$. We denote the core knot of Σ' by K , a meridian of Σ' on $\partial\Sigma'$ by K' , the p' -component link included in Σ by $K_1 \cup \dots \cup K_{p'}$, and the q' -component link included in Σ' by $K'_1 \cup \dots \cup K'_{q'}$. We take $L = K_1 \cup \dots \cup K_{p'} \cup K$ and $L' = K'_1 \cup \dots \cup K'_{q'} \cup K'$. Then $p = p' + 1$, $q = q' + 1$, and $\widehat{L} = S_{pq}(L, L')$. We show that L is a p -component non-split semi-Brunnian link and L' is a q -component non-split Brunnian link.

Both Σ and Σ' include at least one component of $\widehat{L}$. Firstly we show that L is a semi-Brunnian link. Since $\widehat{L}$ is a Brunnian link, $L \setminus K = \widehat{L} \setminus (L' \setminus K')$ is a trivial link. For $i \in \{1, \dots, p'\}$, we suppose that K is a non-split component or a non-trivial component in $L \setminus K_i$. Then a longitude of K does not represent the trivial element in the fundamental group of the complement of $L \setminus (K_i \cup K)$. We note that the longitude of K can be situated on $T = \partial\Sigma$, and we denote it by l . The meridian of Σ' , which is a longitude of K' , is also on $T = \partial\Sigma'$, and we denote it by l' . For the complement of $\widehat{L} \setminus K_i$, since both l and l' are essential in Σ and Σ' respectively, T is still essential. It contradicts that $\widehat{L} \setminus K_i$ is a trivial link. Hence K is an unknot and a split component in $L \setminus K_i$, and $L \setminus K_i$ is a trivial link. Therefore L is a semi-Brunnian link.

Secondly we show that L' is a Brunnian link. Since $\widehat{L}$ is a Brunnian link, $L' \setminus K' = \widehat{L} \setminus (L \setminus K)$ is a trivial link. For $j \in \{1, \dots, q'\}$, we suppose that $L' \setminus K'_j$ is not a trivial link. Then since $L' \setminus (K'_j \cup K')$ is a trivial link, a longitude of K' does not represent the trivial element in the fundamental group of the complement of $L' \setminus (K'_j \cup K')$. For the complement of $\widehat{L} \setminus K'_j$, since l is essential in Σ and l' is essential in Σ' , T is still essential. It contradicts that $\widehat{L} \setminus K'_j$ is a trivial link. Hence $L' \setminus K'_j$ is a trivial link, and L' is a Brunnian link. ■

For the case $p = 1$ and $q \geq 3$, the converse of Lemma 6.9 (3) does not hold.

Example 6.11 Let $R_\lambda = K_1 \cup \dots \cup K_\lambda$ be a λ -component rubberband Brunnian link [Ba] with $\lambda \geq 3$, and $\overline{R}_\lambda = R_\lambda \cup K'$ as in Figure 6, where $K_{\lambda+1} = K'$. Then R_λ is a non-split atoroidal Brunnian link, and $\overline{R}_\lambda$ is not a Brunnian link but an algebraically split link. For a non-trivial knot K , $\widehat{L} = S_{1,\lambda+1}(K, \overline{R}_\lambda)$ is a non-split toroidal Brunnian link.




Figure 6: a λ -component rubberband Brunnian link R_λ and $\overline{R}_\lambda = R_\lambda \cup K'$

Remark 6.12 Since a non-split Brunnian link is prime and anannular, a non-split atoroidal Brunnian link is hyperbolic. There exists infinitely many hyperbolic Brunnian links [Kn]. For example, a rubberband Brunnian link R_λ (see also Example 6.11 and Figure 6) is hyperbolic.

A λ -component link $L = K_1 \cup \dots \cup K_\lambda$ in S^3 with $\lambda \geq 2$ is a *semi-Brunnian type link* if for $\lambda \geq 3$, L is algebraically split link such that for fixed $i \in \{1, 2, \dots, \lambda\}$, K_r ($r \in \{1, 2, \dots, \lambda\} \setminus \{i\}$) is an unknot and the Alexander polynomial of every proper sublink with at least two components is zero, and for $\lambda = 2$ and fixed $i \in \{1, 2\}$, K_r ($r \in \{1, 2\} \setminus \{i\}$) is an unknot. Then K_i is called the *characteristic component* of L .

Theorem 6.13 *Let $L = K_1 \cup \dots \cup K_p$ be a p -component semi-Brunnian type link with $p \geq 2$, K_i the characteristic component of L , and $L' = K'_1 \cup \dots \cup K'_q$ a q -component Brunnian type link with $q \geq 2$. Then $\widehat{L} = S_{ij}(L, L')$ is a $(p+q-2)$ -component Brunnian type link in S^3 for any $j \in \{1, 2, \dots, q\}$.*

To prove Theorem 6.13, we prepare a lemma.

Lemma 6.14 *In the same situation as in Theorem 6.13, we assume that $i = p$ and $j = q$. Then we have the following:*

(1)

$$\Delta_L(t_1, \dots, t_p) \doteq \begin{cases} \frac{(t_1 t_2)^k - 1}{t_1 t_2 - 1} \Delta_{K_2}(t_2) + (t_1 - 1)(t_2 - 1)f(t_1, t_2) & (p = 2), \\ (t_1 - 1) \cdots (t_p - 1)f(t_1, \dots, t_p) & (p \geq 3), \end{cases}$$

where $k = \text{lk}(K_1, K_2)$ for $p = 2$, and

$$\Delta_{L'}(t'_1, \dots, t'_q) \doteq (t'_1 - 1) \cdots (t'_q - 1)f'(t'_1, \dots, t'_q),$$

where $f(t_1, t_2) \in \mathbb{Z}[t_1^{\pm 1}, t_2^{\pm 1}]$, $f(t_1, \dots, t_p) \in \mathbb{Z}[t_1^{\pm 1}, \dots, t_p^{\pm 1}]$, and $f'(t'_1, \dots, t'_q) \in \mathbb{Z}[t'_1, \dots, t'_q]$.

(2) If $p = 2$ and $k \neq 0$, then we have

$$\Delta_{\widehat{L}}(t_1, t'_1, \dots, t'_{q-1}) \doteq \frac{(t_1^k - 1)^2}{t_1 - 1} (t'_1 - 1) \cdots (t'_{q-1} - 1) f'(t'_1, \dots, t'_{q-1}, t_1^k).$$

(3) If $p = 2$ and $k = 0$, or $p \geq 3$, then we have

$$\Delta_{\widehat{L}}(t_1, \dots, t_{p-1}, t'_1, \dots, t'_{q-1}) \doteq 0.$$

Proof. (1) By applying Lemma 2.4 repeatedly, we have the result.

(2) By (1) and Lemma 2.2, we have the result.

(3) We add one component K to L such that $K_i \cup K$ ($i = 1, \dots, p$) is the connected sum of K_i and the Hopf link, where $\text{lk}(K_i, K) = 1$, and we denote by $\overline{L} = L \cup K$ and $\widehat{\overline{L}} = \widehat{L} \cup K$. Then by (1), (3.4) and Lemma 2.2, we have

$$\begin{aligned} \Delta_{\widehat{\overline{L}}}(t_1, \dots, t_{p-1}, t'_1, \dots, t'_{q-1}, t) &\doteq \Delta_{\overline{L}}(t_1, \dots, t_{p-1}, 1, t) \Delta_{L'}(t'_1, \dots, t'_{q-1}, t) \\ &\doteq (t - 1)^2 g(t_1, \dots, t_{p-1}, 1, t) \\ &\quad \times (t'_1 - 1) \cdots (t'_{q-1} - 1) f'(t'_1, \dots, t'_{q-1}, t), \end{aligned}$$

where $g(t_1, \dots, t_{p-1}, 1, t) \in \mathbb{Z}[t_1^{\pm 1}, \dots, t_{p-1}^{\pm 1}, t^{\pm 1}]$. By Lemma 2.4, we have

$$\Delta_{\widehat{\overline{L}}}(t_1, \dots, t_{p-1}, t'_1, \dots, t'_{q-1}, 1) \doteq (t_1 \cdots t_{p-1} - 1) \Delta_{\widehat{L}}(t_1, \dots, t_{p-1}, t'_1, \dots, t'_{q-1}) = 0,$$

and

$$\Delta_{\widehat{L}}(t_1, \dots, t_{p-1}, t'_1, \dots, t'_{q-1}) = 0. \quad \blacksquare$$

Remark 6.15 By Lemma 6.14 (3), we obtain $\Delta_{M_\lambda}(t_1, \dots, t_\lambda) = 0$ for $\lambda \geq 4$ (cf. (1.1)).

Proof of Theorem 6.13. It is easy to see that $\widehat{L}$ is a $(p + q - 2)$ -component link such that every component is an unknot. We show that the Alexander polynomial of every proper sublink of $\widehat{L}$ with at least two components is zero. We suppose that $i = p$ and $j = q$. By Lemma 6.14 (2) and (3), we have the result. $\blacksquare$

Corollary 6.16 Let $L = K_1 \cup K_2$ be a 2-component semi-Brunnian type link, K_2 the characteristic component of L , $L' = K'_1 \cup \cdots \cup K'_q$ a q -component Brunnian type link with $q \geq 2$, and $\widehat{L} = S_{2q}(L, L')$.

(1) If there is another expression of $\widehat{L}$ which is a satellite of a p' -component semi-Brunnian type link with pattern a q' -component Brunnian type link such that $p' \geq 3$ and $q' \geq 2$, then we have

$$\Delta_{\widehat{L}}(t_1, t'_1, \dots, t'_{q-1}) = 0.$$

(2) If $|\text{lk}(K_1, K_2)| \geq 2$ and $\Delta_{L'}(t'_1, \dots, t'_q) \neq 0$, then $\widehat{L}$ does not yield a lens space by any finite surgery.

Proof. (1) By Lemma 6.14 (3), we have the result.

(2) By Lemma 6.14 (2), we have

$$\Delta_{\widehat{L}}(t_1, t'_1, \dots, t'_{q-1}) \doteq \left(\frac{t_1^k - 1}{t_1 - 1} \right)^2 (t_1 - 1)(t'_1 - 1) \cdots (t'_{q-1} - 1) f'(t'_1, \dots, t'_{q-1}, t_1^k),$$

where $k = \text{lk}(K_1, K_2)$. We set

$$f(t_1, t'_1, \dots, t'_{q-1}) = \left(\frac{t_1^k - 1}{t_1 - 1} \right)^2 f'(t'_1, \dots, t'_{q-1}, t_1^k).$$

We prove that a Laurent polynomial which is obtained by substituting 1 to the variables in $f(t_1, t'_1, \dots, t'_{q-1})$ except one variable is not monic or does not satisfy the conditions in Theorem 6.4 and Theorem 6.6. It shows that $\widehat{L}$ does not yield a lens space by any finite surgery by (6.1), Theorem 6.4 and Theorem 6.6.

If we substitute 1 except t'_i to $f(t_1, t'_1, \dots, t'_{q-1})$, then we obtain a Laurent polynomial of the form $k^2 g_i(t'_i)$ where $g_i(t'_i) \in \mathbb{Z}[t'_i, t_i'^{-1}]$. If $|k| \geq 2$, then $f_i(t'_i)$ is not monic. If we substitute 1 except t_1 to $f(t_1, t'_1, \dots, t'_{q-1})$, then we obtain a Laurent polynomial of the form

$$\left(\frac{t_1^k - 1}{t_1 - 1} \right)^2 f'(1, \dots, 1, t_1^k). \quad (6.2)$$

If $|k| \geq 2$, then the trace of the term $\left(\frac{t_1^k - 1}{t_1 - 1} \right)^2$ is -2 and that of the term $f'(1, \dots, 1, t_1^k)$ is 0. Since the degree of (6.2) is at least 2, both the trace of (6.2) and that of $\Delta_{\widehat{K}_1}(t_1)$ in (6.1) are equal to -2 . It does not satisfy the condition in Theorem 6.6. ■

Finally, we pose the following questions :

Question 6.17 (1) Are the Whitehead link and the Borromean rings the only non-trivial Brunnian links and anannular Brunnian type links yielding lens spaces ?

(2) Do every non-trivial Brunnian link and anannular Brunnian type link have Properties FP' , FP and FR ?

Corresponding to the cyclic surgery theorem [CGLS], we ask the following :

Question 6.18 Are the cyclic surgery coefficients of the components of a hyperbolic Brunnian link (resp. a hyperbolic anannular Brunnian type link) integers except that of one component ?

Theorem 1.1, Lemma 3.6 (1) and Theorem 6.7 are partial answers for Question 6.18 above.

Acknowledgements The author would like to thank to Yasuyoshi Tsutsumi for giving him useful advices about the Casson-Walker invariant, Akio Kawauchi for pointing out relationship between the present result and his results [Kw3] from the imitation theory [Kw1, Kw2], and Motoo Tange for informing him about a result of M. Hedden and L. Watson [HW].

References

- [Ba] D. Bar-Natan, “*Rubberband*” *Brunnian link–Knot Atlas*,
<http://katlas.org/wiki/%22Rubberband%22BrunnerianLinks>.
- [CGLS] M. Culler, M. Gordon, J. Luecke and P. Shalen, *Dehn surgery on knots*, Ann. of Math., **125** (1987), 237–300.
- [Gb1] D. Gabai, *The Murasugi sum is a natural geometric operation*, Amer. Math. Soc. Comtemp. Math. **20** (1983), 131–143.
- [Gb2] D. Gabai, *Foliations and the topology of 3-manifolds III*, J. Diff. Geom. **26** (1987), 479–536.
- [GL] C. McA. Gordon and J. Luecke, *Knots are determined by their complements*, J. Amer. Math. Soc. **2** (1989), 371–415.
- [HW] M. Hedden and L. Watson, *On the geography and botany of knot Floer homology*, arXiv: math.GT/1404.6913.
- [Kd1] T. Kadokami, *Reidemeister torsion of Seifert fibered homology lens spaces and Dehn surgery*, Algebr. Geom. Topol. **7** (2007), 1509–1529.
- [Kd2] T. Kadokami, *Reidemeister torsion and lens surgeries on knots in homology 3-spheres II*, Top. Appl. **155**, No.15 (2008), 1699–1707.
- [Kd3] T. Kadokami, *Amphicheiral links with special properties, I*, J. Knot Theory Ramif. **21** No. 6 (2012) 1250048, 17p.
- [KMS] T. Kadokami, N. Maruyama and M. Shimoizawa, *Lens surgeries along the n -twisted Whitehead link*, Kyungpook Math. J. **52** No. 3 (2012), 245–264.
- [KS] T. Kadokami and M. Shimoizawa, *Dehn surgery along torus links*, J. Knot Theory Ramif. **19** No. 4 (2010), 489–502.
- [KY] T. Kadokami and Y. Yamada, *A deformation of the Alexander polynomials of knots yielding lens spaces*, Bull. of Austral. Math. Soc. **75** (2007), 75–89.
- [Kn] T. Kanenobu, *Hyperbolic links with Brunnian properties*, J. Math. Soc. Japan **38** No.2 (1986), 295–308.
- [Kw1] A. Kawauchi, *An imitation theory of manifolds*, Osaka J. Math. **26** (1989), 447–464.
- [Kw2] A. Kawauchi, *Topological imitation of a colored link with the same Dehn surgery manifold*, Top. Appl. **146–147** (2005), 67–82.
- [Kw3] A. Kawauchi, *Recent applications of topological imitation theory*, talk in East China Normal University (2011).

- [KM] P. Kronheimer and T. Mrowka, *Witten's conjecture and Property P*, Geom. Topol., **8** (2004), 295–310.
- [KMOS] P. Kronheimer, T. Mrowka, P. Ozsváth and Z. Szabó, *Monopoles and lens space surgeries*, Ann. of Math. (2), **165** No. 2 (2007), 457–546.
- [Les] C. Lescop, *Global surgery formula for the Casson-Walker invariant*, Ann. of Math. Studies, Princeton Univ. Press. **140** (1996).
- [MS] B. Mangum and T. Stanford, *Brunnian links are determined by their complements*, Algebr. Geom. Topol. **1** (2001), 143–152.
- [MP] B. Martelli and C. Petronio, *Dehn filling of the “magic” 3-manifold*, Comm. Anal. Geom. **14** No. 5 (2006), 969–1026.
- [Mi1] J. Milnor, *Link groups*, Ann. of Math. (2) **59** (1954), 177–195.
- [Mi2] J. Milnor, *Isotopy of links*, Algebraic geometry and topology, A symposium in honor of S. Lefschetz, pp. 280–306, Princeton University Press, Princeton, N. J., 1957.
- [Mi3] J. W. Milnor, *A duality theorem for Reidemeister torsion*, Ann. of Math. (2) **76** (1962), 137–147.
- [Mos] L. Moser, *Elementary surgery along a torus knot*, Pacific J. Math. **38** (1971), 737–745.
- [Ni] Y. Ni, *Knot Floer homology detects fibered knots*, Invent. Math. **170** (2007), 577–608.
- [Nc] L. Nicolaescu, *Seiberg-Witten invariants of rational homology 3-spheres*, Commun. Contemp. Math., **6** No. 6 (2004), 833–866.
- [OS] P. Ozsváth and Z. Szabó, *On knot Floer homology and lens space surgeries*, Topology, **44** (2005), 1281–1300.
- [Pe1] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, arXiv: math.DG/0211159.
- [Pe2] G. Perelman, *Ricci flow with surgery on three-manifolds*, arXiv: math.DG/0303109.
- [Pe3] G. Perelman, *Finite extinction time for the solutions to the Ricci flow on certain three-manifolds*, arXiv: math.DG/0307245.
- [Ro] D. Rolfsen, *Rational surgery calculus: extension of Kirby's theorem*, Pacific J. Math. **110** (1984), 377–386.
- [St] T. Sakai, *Reidemeister torsion of a homology lens space*, Kobe J. Math. **1** (1984), 47–50.
- [Ta] M. Tange, *On the Alexander polynomial of lens space knot*, arXiv: math.GT/1409.7032.
- [To] G. Torres, *On the Alexander polynomial*, Ann. of Math. (2) **57** (1953), 57–89.
- [Tr1] V. G. Turaev, *Reidemeister torsion in knot theory*, Russian Math. Surveys **41-1** (1986), 119–182.
- [Tr2] V. G. Turaev, *Introduction to Combinatorial Torsions*, Birkhäuser Verlag (2001).
- [Wa] K. Walker, *An extension of Casson's invariant*, Ann. of Math. Stud. **126**, Princeton University Press, (1992).
- [Was] L. C. Washington, *Introduction to Cyclotomic Fields*, Graduate Texts in Mathematics **83**, Springer-Verlag, New York (1982).

Teruhisa KADOKAMI
 Department of Mathematics, East China Normal University,
 Dongchuan-lu 500, Shanghai, 200241, China
 mshj@math.ecnu.edu.cn
 kadokami2007@yahoo.co.jp